



PollenX™

FOLIAR SPRAY TO SUPPORT FLOWER AND SET.

GUARANTEED ANALYSIS

Available Phosphate (P ₂ O ₅)	1.10 %
Calcium (Ca)	0.50 %

DENSITY: 8.72 LBS/GAL @ 68°F (20°C)
STORE ABOVE 40°F

Derived From: Calcium Chloride, Potassium Phosphate Chlorine (Cl), Maximum not more than 1.00%

ALSO CONTAINS NONPLANT FOOD INGREDIENT:
87% Kelp Extract (*Ecklonia maxima*) (Microbe Food)

KEEP OUT OF REACH OF CHILDREN.

This product does not meet the hazardous criteria set forth by the 2012 OSHA Hazard Communication Standard (29 CFR 1910.1200). However, observe good industrial hygiene practices. Do not breathe spray. Wash face, hands and skin thoroughly after handling. Wear protective gloves, protective clothing and eye protection. Wash contaminated clothing before reuse. IF SWALLOWED: Rinse mouth. IF ON SKIN (or hair): Immediately remove all contaminated clothing. Rinse skin with water. IF INHALED: If breathing is difficult, move to fresh air and rest in a position comfortable for breathing. IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing.



Technovation

Specialized to succeed

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.aapfc.org/metals.html>

331 Almahurst Lane
Nicholasville
Kentucky
info@agritechnovation.com
www.agritechnovationinc.com

NET CONTENTS:
2.5 GALLONS (9.5 LITERS)
5 GALLONS (18.9 LITERS)

NET WEIGHT
21.8 LBS (12.11 KG)
43.6 LBS (19.77 KG)

DIRECTIONS FOR USE

BRILLIANCE™ is recommended for use on all crops during reproductive growth phases to support development. BRILLIANCE™ is fully water soluble and provides readily available nutrients as shown in the guaranteed analysis for foliar applications. BRILLIANCE™ will be influenced by differences in varieties, soil, and environmental conditions and should be applied under conditions of optimum plant growth. Optimum conditions are defined as adequate humidity, moderate temperatures, and adequate soil moisture. Optimum conditions for foliar application to most crops generally occur during the early morning or late evening hours.

- Avoid application under extreme conditions such as temperatures above 86°F and wind speeds exceeding 5 mph.
- Wear suitable protective clothing when sprayed with other products.
- Optimum results with this product are obtained if the pH of the tank mix is between 5 and 6.
- This product should be applied as a foliar spray during phenological stages indicated in the application table.
- For optimum results BRILLIANCE™ should be used in a plant nutrition program. Recommendations should be based on soil and leaf analyses.
- Where overhead irrigation systems are in use, avoid scheduling irrigation directly after application.

CAUTION

Do not apply when plants are under water or heat stress, or during cool wet conditions. Do not store mixed product for more than 8 hours. Do NOT mix or apply in combination with amine herbicides, copper containing fungicides or phosphonates. Do not exceed the appropriate application rates. Consult with your distributor, crop adviser or manufacturer should crop specific programs, or any other information be required around the use of this product.

STORAGE & DISPOSAL:

Store product in original container. DO NOT store in excessive cold or heat or direct sunlight. DO NOT re-use container. DO NOT contaminate water, food or feed by storage or disposal. Keep packaging closed, securely fastened and upright. Triple rinse or equivalent and dispose of container in accordance with local, state, and federal regulations.

APPLICATION RATES

The rate of application is crop dependent. Apply BRILLIANCE™ according to label rates. For other modes of application, consult a qualified advisor.

MIXING

Agitate BRILLIANCE™ well before dilution or use. Always mix BRILLIANCE™ with sufficient quantities of water. Never apply BRILLIANCE™ undiluted. Follow standard tank mixing procedures. Ensure good agitation throughout the application.

COMPATIBILITY

It is advisable to do a miscibility test prior to mixing with any other product and to consult with a qualified advisor or distributor.

DISCLAIMER

The manufacturer, importer and seller make no express or implied warranties, including any warranty of merchantability, fitness for a particular purpose, or suitability. The product is sold "AS-IS". All sales are final. In no event will manufacturer, importer and seller have any liability for any damages, whether direct, incidental, special or consequential arising out of or in connection with any duty with respect to the product, including, but not limited to lost profits, lost sales or injury to persons or property. Manufacturer, importer and seller's liability on any claim of any kind, including but not limited to warranty, negligence, strict liability, and any other cause of action, for any loss or damage arising out of, connected with, or resulting from the performance or breach of the terms of sale of any goods, or any related services related to sale, delivery, resale, technical direction, inspection, operation or use of any goods or services or part of those goods or services will in no case exceed the purchase price allocable to the goods or services or part of those goods or services that gives rise to the claim. Manufacturer, importer and seller make no representation whatsoever with regard to whether any goods comply with the Occupational Safety and Health Act (OSHA), the Americans with Disabilities Act (ADA), or any other federal, state or local statute, law, ordinance or ruling. Customer, buyer, and user assume all responsibility for use of product.

APPLICATION RATES

Crop	Maximum Rate per Acre*	Rate per 100 gal Water	Additional Information
Corn	1 to 3 qt	40 - 60 fl oz	Apply as a foliar at V12 until silking
Canola	1 to 3 qt	40 - 60 fl oz	Apply as a foliar at flowering
Soybeans	1 to 3 qt	40 - 60 fl oz	Apply as a foliar at R1/R2
Wheat/barley	1 to 2 qt	40 - 50 fl oz	Apply as a foliar at BBCH 60 to 69 stages

*Maximum rates are dependent on crop size.