



COPPER-TO-PERFORM™

F002806

LIQUID COPPER FOR FOLIAR APPLICATION

GUARANTEED ANALYSIS

Sulfur (S)	2.00%
2.0% Combined Sulfur (S)	
Copper (Cu)	5.40%
5.4% Water Soluble Copper (Cu)	

Derived From: Copper Sulfate

DENSITY: 9.85 LBS/GAL @ 68°F (20°C)

STORE ABOVE 40°F

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.aapfco.org/metals.html>



KEEP OUT OF REACH OF CHILDREN.

DANGER Causes serious eye irritation. Harmful if swallowed.

Wash face, hands and skin thoroughly after handling. Wear eye protection. Do not eat, drink or smoke when using this product. IF SWALLOWED: Rinse mouth. Immediately call a POISON CENTER or doctor. IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. Immediately call a POISON CENTER or doctor. Dispose of container in accordance with regulations.



Technovation

Specialized to succeed

**NET CONTENTS: 2.5 Gallons (9.5 Liters)
5 Gallons (18.9 Liters)**

**NET WEIGHT: 24.6 LBS (11.2 KG)
49.3 LBS (22.4 KG)**

331 Almahurst Lane
Nicholasville
Kentucky
40356

DIRECTIONS FOR USE

COPPER-TO-PERFORM™ is a liquid, effective, readily available source of copper and sulfur for soil and foliar applications.

COPPER-TO-PERFORM™ will be influenced by differences in crop varieties, soil and environmental conditions and should be applied under conditions of optimum plant growth. Optimum conditions are defined as adequate humidity, moderate temperatures and adequate soil moisture. Optimum conditions for foliar application to most crops generally occur during early morning or late evening hours.

- Do not apply to crops that are water or heat stressed or in very windy conditions.
- Wear suitable protective clothing when sprayed with other products.
- For best results with this product, the pH of the tank mixture should be between 5 and 6.
- This product should be applied as a foliar spray when crops have sufficient leaf area.
- For optimum results COPPER-TO-PERFORM™ should be used in a plant nutrition program. Recommendations should be based on soil and leaf analyses.
- Where overhead systems are in use, avoid scheduling irrigation directly after application.
- In orchards equipped with micro irrigation systems, spray deposits should not be affected by irrigation.

CAUTION

Do not apply on wet fruit. Application can only take place once the fruit is dry. Do not apply to plants that are undergoing a period of moisture or heat stress, or during cool wet conditions.

STORAGE & DISPOSAL

Store product in original container. DO NOT store in excessive cold or heat or direct sunlight. DO NOT reuse container. DO NOT contaminate water, food or feed by storage or disposal. Keep packaging closed, securely fastened and upright. Triple rinse or equivalent and dispose of container in accordance with local, state and federal regulations. Do not exceed the appropriate application rates.

APPLICATION RATES

The rate of application is dependent on the soil fertility and reaction required. For more information on foliar application, see table on application rates. For other modes of application, consult a qualified advisor.

MIXING

Shake COPPER-TO-PERFORM™ well before dilution or use. Always mix COPPER-TO-PERFORM™ with sufficient quantities of water. Never apply COPPER-TO-PERFORM™ undiluted. Follow standard tank mixing procedures. Ensure good agitation throughout application.

COMPATIBILITY

It is advisable to do a miscibility test prior to mixing with any other product and to consult with a qualified advisor or distributor. Do not spray COPPER-TO-PERFORM™ within 30 days after application of any Copper containing products. Do not store mixed product for more than 8 hours. Do NOT mix or apply in combination with Calcium containing products, amine herbicides, Copper containing fungicides, Sulfates, Phosphates, phosphonates, highly alkaline material such as Lime Sulfur, MCPA or highly alkaline Boron products

DISCLAIMER

The manufacturer, importer and seller make no express or implied warranties, including any warranty of merchantability, fitness for a particular purpose, or suitability. The product is sold "AS-IS". All sales are final. In no event will manufacturer, importer and seller have any liability for any damages, whether direct, incidental, special or consequential arising out of or in connection with any duty with respect to the product, including, but not limited to lost profits, lost sales or injury to persons or property. Manufacturer, importer and seller's liability on any claim of any kind, including but not limited to warranty, negligence, strict liability, and any other cause of action, for any loss or damage arising out of, connected with, or resulting from the performance or breach of the terms of sale of any goods, or any related services related to sale, delivery, resale, technical direction, inspection, operation or use of any goods or services or part of those goods or services will in no case exceed the

purchase price allocable to the goods or services or part of those goods or services that gives rise to the claim. Manufacturer, importer and seller make no representation whatsoever with regard to whether any goods comply with the Occupational Safety and Health Act (OSHA), the Americans with Disabilities Act (ADA), or any other federal, state or local statute, law, ordinance or ruling. Customer, buyer and user assume all responsibility for use of product.

APPLICATION RATES

Crop	Maximum Rate per Acre*	Rate per 100 gal Water	Additional Information
Citrus	2.5 qt	1 pt	First application at green bud. Repeat at 100% petal fall and again at cell enlargement stage.
Grapes	2.5 pt	1 - 2.5 pt	Start first application at 3-6 leaf phase and again at 20 inch shoot growth.
Corn, Cotton	5 fl. oz.	1 pt	Apply early in season between 6-8 leaf stage. Repeat sprays if required.
Macadamias and other nuts	1 - 2 qt	1 pt	First application on first Spring flush, second application after flower and again at cell enlargement stage.
Pastures	7 fl. oz.	1 pt	Apply at 2-4 leaf stage and later new growth.
Pecan nuts	1 - 2 qt	1 pt	Start first application in spring after budbreak and on all new growth in the summer. Three to six applications are recommended depending on requirements
Pome and Stone Fruit	3 - 4 pt	10 - 13 fl. oz.	Apply as full cover spray at bud burst and repeat 2-3 times depending on requirements, every two weeks after petal drop. Apply one more application after harvest. Do not spray in flower time.
Soybeans, Green Beans and Peanuts	1 - 2 qt	1 pt	Apply early in season between 6-8 leaf stage. Repeat sprays if required.
Tomatoes, Chillies and Peppers	1 qt	2 - 3 qt	Apply at 2-4 leaf stage Repeat 3 more times after in 10-day intervals.
Wheat, Oats, Canola and Barley	5 fl. oz.	1 pt	Apply at 2-4 leaf stage. Repeat if necessary.
Almonds	1 - 2 qt	1 pt	Apply before flowering on spring flush repeat just after flowering and again in the fall.
Pistachios	1 - 2 qt	1 pt	Apply on new spring flush repeat just after flowering and again in the fall.

*Maximum rate depends on crop size.