



K-TO-PERFORM™

F002806

LIQUID POTASSIUM FOR FOLIAR APPLICATION

0 - 0 - 23

GUARANTEED ANALYSIS

Soluble Potash (K ₂ O)	23.00%
-----------------------------------	--------

Derived from: Potassium Citrate, Potassium Acetate

DENSITY: 10.51 LBS/GAL @ 68°F (20°C)

STORE ABOVE 40°F

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.aapfco.org/metals.html>



KEEP OUT OF REACH OF CHILDREN.

This product does not meet the hazardous criteria set forth by the 2012 OSHA Hazard Communication Standard (29 CFR 1910.1200). However, observe good industrial hygiene practices. Do not breathe spray. Wash face, hands and skin thoroughly after handling. Wear protective gloves, protective clothing and eye protection. Wash contaminated clothing before reuse. IF SWALLOWED: Rinse mouth. IF ON SKIN (or hair): Immediately remove all contaminated clothing. Rinse skin with water. IF INHALED: If breathing is difficult, move to fresh air and rest in a position comfortable for breathing. IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing.



Technovation

Specialized to succeed

**NET CONTENTS: 5 Gallons (18.9 Liters)
2.5 Gallons (9.5 Liters)**

**NET WEIGHT: 52.6 LBS (23.8 KG)
26.3 LBS (11.9 KG)**

331 Almahurst Lane
Nicholasville
Kentucky
40356

DIRECTIONS FOR USE

K-TO-PERFORM™ is recommended for use on all crops during plant growth periods that require relatively high potassium levels, such as during flowering and fruit formation when additional nitrogen will be detrimental to the crop quality. K-TO-PERFORM™ is completely water soluble.

K-To-PERFORM™ is low in chlorine with a maximum chlorine (Cl) content of 0.10%.

K-TO-PERFORM™ should preferably be applied in the early morning or late afternoon.

- Keep out of direct sunlight.
- Keep container closed when not in use.

CAUTION

Do not apply to crops that are water or heat stressed. Do not spray K-TO-PERFORM™ within 21 days after copper application. Do NOT tank mix K-TO-PERFORM™ with copper-containing products.

Consult with your distributor, crop adviser or manufacturer should crop specific programs, or any other information be required around the use of this product.

APPLICATION RATES

The rate of application is dependent on the soil fertility and reaction required. See table of recommended application rates. Lower rates should be used for maintenance while higher rates will correct nutritional deficiencies.

MIXING

Always mix K-TO-PERFORM™ with sufficient quantities of water. Never apply K-TO-PERFORM™ undiluted. Follow standard tank mixing procedures. Ensure good agitation throughout application.

COMPATIBILITY

It is advisable to do a miscibility test prior to mixing with any other product and to consult with a qualified advisor or distributor.

DISCLAIMER

The manufacturer, importer and seller make no express or implied warranties, including any warranty of merchantability, fitness for a particular purpose, or suitability. The product is sold "AS-IS". All sales are final. In no event will manufacturer, importer and seller have any liability for any damages, whether direct, incidental, special or consequential arising out of or in connection with any duty with respect to the product, including, but not limited to lost profits, lost sales or injury to persons or property. Manufacturer, importer and seller's liability on any claim of any kind, including but not limited to warranty, negligence, strict liability, and any other cause of action, for any loss or damage arising out of, connected with, or resulting from the performance or breach of the terms of sale of any goods, or any related services related to sale, delivery, resale, technical direction, inspection, operation or use of any goods or services or part of those goods or services will in no case exceed the purchase price allocable to the goods or services or part of those goods or services that gives rise to the claim. Manufacturer, importer and seller make no representation whatsoever with regard to whether any goods comply with the Occupational Safety and Health Act (OSHA), the Americans with Disabilities Act (ADA), or any other federal, state or local statute, law, ordinance or ruling. Customer, buyer and user assume all responsibility for use of product.

APPLICATION RATES

Application	Maximum Rate per Acre*	Rate per 100 gallons of Water	Additional Information
Fruit and Nuts	1 - 3 qt	2 qt	Start application as soon as the cell enlargement stage begins or Potassium levels become limiting. Apply as a full cover spray. Repeat at 1 to 2 week intervals as required.
Grains and Sugarcane	1 - 2 qt	2 - 3 qt	Start application as soon as the cell enlargement stage begins or Potassium levels become limiting. Apply as a full cover spray. Repeat at 1 to 2 week intervals as required.
Vegetables	1 - 2 qt	2 - 3 qt	Start application as soon as the cell enlargement stage begins or Potassium levels become limiting. Apply as a full cover spray. Repeat at 1 to 2 week intervals as required.
Application	Maximum Rate per Acre	Rate per 5 gallons of water per Acre	Additional Information
Aerial application	3 gallons	2 gallons	Start application as soon as the cell enlargement stage begins or Potassium levels become limiting. Repeat at 1 to 2 week intervals as required.

*Maximum rates depend on crop age.